## THESIS TOPIC

Subject N° (to be completed by the ED):	FUNDING:	⊠ Requested ☐ Acquired	Funding origin:
Thesis title: Functional redundancy within bacterial regulators: characterization of <i>Staphylococcus aureus</i> sRNA co-regulation by transcription factors and involvement in virulence.		3 keywords : S. aureus, Heterologous one hybrid system, sRNA	
Research Unit / team : U1230_BRM, EA2311 / laboratoire de Biochimie pharmaceutique			
Director's name: Astrid Jouvante Rouillon			Phone number: 02 23 23 43 94 Email address: astrid.rouillon@univ-rennes1.fr Year of HDR: 2020
Socio-economic and scientific context (approximately 10 lines): Multidrug resistant bacterial infections (MDR) represent major public health issues and an economic burden. In 2019, 1.2 million deaths were attributed to BMR infection (Murray et al, 2022). These bacteria are a threat because of the difficulty in finding effective treatments for infections. Among ESKAPE, <i>S. aureus</i> is an asymptomatically borne commensal bacterium in about 30% of the population. It is involved in more than half of skin infections but is also responsible for bacteremia, osteomyelitis and endocarditis. The success of an <i>S. aureus</i> infection is linked to its ability to coordinate the expression of many virulence factors at key moments, but also to adapt to environmental changes. To this end, <i>S. aureus</i> expresses protein regulators such as transcription factors (TF) and 2-component systems (TCS), as well as RNA-type regulators such as RNA regulators (sRNA). Despite a crucial role in transcriptional and post-transcriptional regulation, corresponding genes are rarely essential to the survival of the bacteria. This is because TF and sRNA are embedded in complex regulations where several FT regulate the same sRNA, and where several sRNA regulate the same target (Menard et al, 2022).			
Working hypothesis and aims (approximately 8 lines): These redundancies appear to be a determining factor explaining the absence of essentiality, as well as we expect that the joint absence of a couple or pool of regulators could lead to stronger phenotypes (Rachwalski et al, 2022). Our previous work has focused on this in the study of the direct targets of the TF SarA, in which we showed that 11 of the 37 sRNAs repressed by SarA were also repressed by another TF, CodY (Augagneur et al, 2020; Oriol et al)). Functional redundancy would mask the importance of some regulators. The objective of this thesis project is therefore to reveal the redundancies between sRNA through the identification of common regulators. These will be revealed thanks to the one hydrid system in yeast. Then, we will study the phenotypes related to the joint absence or overexpression of sRNA in particular in virulence, antibiotic resistance and biofilm formation. We will work with a series of about 12 sRNAs, which will have been selected because targets and biological functions have already been associated with them.			
Main milestones of the thesis (approximately 12 lines):			
We will use the heterologous one hybrid system in the yeast Saccharomyces cerevisiae as a screening tool to identify the S. aureus TF regulating the selected S. aureus sRNAs. In this context, a 75 TF expression bank of S. aureus will be built. Similarly, the 12 strains of the carrier yeast for which the 12 regions promoting the sRNAs of interest have been cloned upstream of the reporter gene will be built. Screening will allow the identification of all TF regulating each sRNA of interest. Next, we will select the most promising regulatory duos (sRNA or TF) in terms of functional redundancy to build double mutants in S. aureus. These mutants will be studied in relation to their phenotype in virulence, antibiotic resistance and biofilm formation.			
For the study of <i>in vivo</i> virulence, we opted for the larval model <i>Galleria mellonella</i> , which is already used in the laboratory and is ethically more responsible (Ménard et al, 2021). However, this model requires the development and/or adaptation of tools. This is why, in parallel with the development of screening tools by heterologous one hybrid, it will be necessary to test and choose the tools allowing bacterial visualization and quantification (bioluminescence or fluorescence) during infection. This will be done through the integration of reporter genes into a neutral locus of the bacterial chromosome. This system built in a wild strain will be transferred by phage transduction into the mutants we study. The identification of regulatory duos essential for growth, virulence, biofilm formation or the development of antibiotic resistance could, in the longer term, lead to the identification of antibacterial targets.			
Scientific and technical skills required by the candidate (2 lines): The candidate will need to have a good knowledge of molecular biology as well as microbiology. He/she will need to feel able to perform experiments on the <i>Galleria mellonella</i> larval model.			
<ol> <li><u>3 publications from the team related to the topic (last 5 years)</u>:</li> <li>1. Charlotte Oriol, Liviu Cengher, Adhar C. Manna, Tony Mauro, Marie-Laure Pinel-Marie, Brice Felden, Ambrose Cheung, Astrid Rouillon. Expanding the <i>Staphylococcus aureus</i> SarA Regulon to Small RNAs. mSystems. 2021 Sep-Oct; 6(5): e00713-21.</li> <li>2. Guillaume Ménard, Astrid Rouillon, Gevorg Ghukasyan, Mathieu Emily, Brice Felden, Pierre-Yves Donnio. <i>Galleria mellonella</i> Larvae as an Infection Model to Investigate sRNA-Mediated Pathogenesis in <i>Staphylococcus aureus</i>. Front Cell Infect Microbiol. 2021; 11: 631710.</li> <li>3. Amiot N., Augagneur Y, Camberlein E., Nicolas I., Lecureur V., Rouillon A. et Felden B. A novel <i>Staphylococcus aureus</i> cis-trans type I toxin- antitoxin module with dual effects on bacteria and host cells. Nucleic Acids Res. 2019 Feb 28;47(4):1759-1773.</li> </ol>			
National and international collaborations: C. Nielsen-Leroux, INRAe, MICALIS : teams GME and MicrobAdapt, Jouy en Josas, France			